

Sol.
$$Na_2SO_3 \xrightarrow{\text{dil. } H_2SO_4} SO_2 \xrightarrow{\text{NaOH}} Na_2SO_3$$

(X) (Y) (X) $\downarrow^{SO_2}_{H_2O}$
NaHSO₃ (Z)

6. The IUPAC name of the following compound is :



- (1) 3-amino-4-hydroxymethyl-5-nitrobenzaldehyde
- (2) 2-nitro-4-hydroxymethyl-5-aminobenzaldehyde

(3) 4-amino-2-formyl-5-hydroxymethylnitrobenzene

(4) 5-amino-4-hydroxymethyl-2-nitrobenzaldehyde

Official Ans. by NTA (4)



5-amino-4-hydroxymethyl-2-nitrobenzaldehyde

- Dihydrogen of high purity (> 99.95%) is obtained through:
 - the electrolysis of warm Ba(OH)₂ solution using Ni electrodes.
 - (2) the reaction of Zn with dilute HCl
 - (3) the electrolysis of brine solution.
 - (4) the electrolysis of acidified water using Pt electrodes.

Official Ans. by NTA (1)

Sol. High purity (>99.95%) dihydrogen is obtained by electrolysing warm aqueous barium hydroxide solution between nickel electrodes.

8.	Match the following :						
		Test/Method		Reagent			
	(i)	Lucas Test	(a)	C ₆ H ₅ SO ₂ Cl/aq. KOH			
	(ii)	Dumas method	(b)	HNO ₃ /AgNO ₃			
	(iii)	Kjeldahl's metho	d (c)	CuO/CO ₂			
	(iv)	Hinsberg Test	(d)	Conc. HCl and ZnCl ₂			
			(e)	H_2SO_4			
	(1) (i)-(d), (ii)-(c), (iii)-(e), (iv)-(a)						
	(2) (i)-(b), (ii)-(d), (iii)-(e), (iv)-(a)						
	(3) (i)-(d), (ii)-(c), (iii)-(b), (iv)-(e)						
	(4) (i)-(b), (ii)-(a), (iii)-(c), (iv)-(d) Official Ans. by NTA (1)						
Sol.		Test	Co	orrect reagent			
	(i)	Lucas test \longrightarrow	co	nc. HCl + $ZnCl_2$			
	(ii) Dumas method \longrightarrow CuO / CO ₂						
	(iii) Kjeldahl's method $\longrightarrow H_2SO_4$						
	(iv)	(iv) Hinsberg Test $\longrightarrow C_6H_5SO_2Cl + aq. KOH$					
9.	The reaction of NO with N_2O_4 at 250 K gives						
	(1)	N_2O_5	(2)) NO ₂			
	(3) N_2O (4) N_2O_3						
	Off	icial Ans. by NTA	A (4)				
Sol.	2No	$O + N_2O_4 \xrightarrow{250K}$	$2N_{2}C$),			
001		2 .	blue so	lid			
10.	For	the given cell;					
	$Cu(s) Cu^{2+}(C_1M) Cu^{2+}(C_2M) Cu(s)$ change in						
	Gibbs energy (ΔG) is negative, if :						
	(1)	$C_1 = 2C_2$	(2)	$C_2 = \frac{C1}{\sqrt{2}}$			
				/ \2			
	(3)	$C_1 = C_2$	(4)) $C_2 = \sqrt{2}C_1$			
	Off	icial Ans. by NTA	(4)				
Sol.	$\Delta G = -n \ F \ E_{cell}$ $\Delta G \ is negative, \ if \ E_{cell} \ is \ positive$ Anode : $Cu(s) \longrightarrow Cu^{+2}(C_1) + 2e^- : E^\circ$						
	Cathode : $Cu^{+2}(C_2) + 2e^{-} \longrightarrow Cu(S) : -E^{\circ}$						
	Cell r	reaction : $Cu^{+2}(C_2)$ —	$\rightarrow Cu^{+}$	$^{2}(C_{1}) E_{cell}^{\circ} = 0$			
	$E_{cell} = E_{cell}^{\circ} - \frac{2.303 \text{RT}}{\text{nF}} \log Q$ $E_{cell} = 0 - \frac{2.303 \text{RT}}{\text{nF}} \log \left(\frac{C_1}{C_2}\right)$						
	$E_{cell} > 0: if \frac{C_1}{C_2} < 1 \Longrightarrow C_1 < C_2$						

2

11. A crystal is made up of metal ions $'M_1'$ ana $'M_2'$ and oxide ions. Oxide ions form a ccp lattice structure. The cation $'M_1'$ occupies 50% of octahedral voids and the cation $'M_2'$ occupies 12.5% of tetrahedral voids of oxide lattice. The oxidation numbers of $'M_1'$ and $'M_2'$ are, respectively :

$$(1) + 2, + 4$$
 $(2) + 3, + 1$

Official Ans. by NTA (1)

Sol. O⁻² ions form ccp. $O_4 \downarrow (-8 \text{ charge})$

$$M_1 = 50\% \text{ of O.V.} \Rightarrow \frac{50}{100} \times 4 = 2:(M_1)_2$$

$$M_2 = 12.5\%$$
 of T.V. $\Rightarrow \frac{12.5}{100} \times 8 = 1:(M_2)_1$

So formula is : $(M_1)_2 (M_2)_1 O_4$

This must be neutral. Both metals must have +8 charge in total.

From given options : $\left\{ \begin{array}{c} \text{O.N. of } M_1 = +2 \\ M_2 = +4 \end{array} \right\}$

- **12.** For a d⁴ metal ion in an octahedral field, the correct electronic configuration is :
 - (1) $t_{2g}^4 e_g^0$ when $\Delta_0 < P$
 - (2) $e_g^2 t_{2g}^2$ when $\Delta_0 < P$
 - (3) $t_{2g}^3 e_g^1$ when $\Delta_0 < P$
 - (4) $t_{2\sigma}^3 e_{\sigma}^1$ when $\Delta_0 > P$

Official Ans. by NTA (3)



back pairing is not possible because pairing energy $> \Delta_{O}.$

13. Which of the following compounds can be prepared in good yield by Gabriel phthalimide synthesis?



Official Ans. by NTA (1)

Sol. Gabriel phthalimide synthesis is used for preparation of 1° Aliphatic amine



Here
$$R-Br = \bigcup_{i=1}^{CR}$$

$$R-NH_2 = OCCH_2-NH_2$$

14. The correct match between Item-I (starting material) and Item-II (reagent) for the preparation of benzaldehyde is :

Item-I	Item-II
(I)Benzene	(P)HCl and $SnCl_2$, H_3O^+
(II)Benzonitrile	(Q) H ₂ , Pd-BaSO ₄ , S and quinoline

(III)Benzoyl Chloride (R)CO, HCl and AlCl₃

- (1) (I)-(Q), (II)-(R) and (III)-(P)
- (2) (I)-(R), (II)-(Q) and (III)-(P)
- (3) (I)-(R), (II)-(P) and (III)-(Q)
- (4) (I)-(P), (II)-(Q) and (III)-(R)

Official Ans. by NTA (3)



- 15. The average molar mass of chlorine is 35.5 g mol⁻¹. The ratio of ³⁵Cl to ³⁷Cl in naturally occurring chlorine is close to :
 - (1) 4 : 1
 - (2) 1 : 1
 - (3) 2 : 1
 - (4) 3 : 1
 - Official Ans. by NTA (4)
- Sol. let x : 1 mole ratio $x^{35}Cl = 3^{37}Cl$ Av. molar mass = 35.5

Av. molar mass =
$$\frac{n_1 M_1 + n_2 M_2}{(n_1 + n_2)}$$

$$35.5 = \frac{x \times 35 + 1 \times 37}{x + 1}$$

- **16.** Which one of the following statements not true ?
 - (1) Lactose contains α -glycosidic linkage between C₁ of galactose and C₄ of glucose.
 - (2) Lactose $(C_{11}H_{22}O_{11})$ is a disaccharide and it contains 8 hydroxyl groups.
 - (3) On acid hydrolysis, lactose gives one molecule of D(+)-glucose and one molecule of D(+)-galactose.
 - (4) Lactose is a reducing sugar and it gives Fehling's test.

Official Ans. by NTA (1)

structure of lactose

17. A set of solutions is prepared using 180 g of water as a solvent and 10 g of different non-volatile solutes A, B and C. The relative lowering of vapour pressure in the presence of these solutes are in the order [Given, molar mass of A = 100 g mol⁻¹; B = 200 g mol⁻¹; C = 10,000 g mol⁻¹]

Sol. Relative lowering of V.P. =
$$\frac{\Delta P}{P^0} = x_{solute}$$

$$\left(\frac{\Delta P}{P^0}\right)_A = \frac{\frac{10}{100}}{\frac{10}{100} + \frac{180}{18}} : \left(\frac{\Delta P}{P^0}\right)_B = \frac{\frac{10}{200}}{\frac{10}{200} + \frac{180}{18}}$$

$$\left(\frac{\Delta P}{P^0}\right)_{C} = \frac{\frac{10}{10,000}}{\frac{10}{10,000} + \frac{180}{18}} : \left(\frac{\Delta P}{P^0}\right)_{A} > \left(\frac{\Delta P}{P^0}\right)_{B} > \left(\frac{\Delta P}{P^0}\right)_{C}$$

18. For a reaction,

10

 $\label{eq:4M(s) + nO_2(g) } \textbf{ } \rightarrow 2M_2O_n(s),$

the free energy change is plotted as a function of temperature. The temperature below which the oxide is stable could be inferred from the plot as the point at which :

- (1) the slope changes from positive to zero
- (2) the free energy change shows a change from negative to positive value
- (3) the slope changes from negative to positive(4) the slope changes from positive to negativeOfficial Ans. by NTA (2)

19. Match the following compounds (Column-I) with their uses (Column-II) :

S.No.	Column – I	S.No.	Column – II
(I)	Ca(OH) ₂	(A)	casts of statues
(II)	NaCl	(B)	white wash
(III)	$CaSO_4 \cdot \frac{1}{2}H_2O$	(C)	antacid
	CaCO ₃	(D)	washing soda
(1V)			preparation

- (1) (I)-(D), (II)-(A), (III)-(C), (IV)-(B)
- (2) (I)-(B), (II)-(C), (III)-(D), (IV)-(A)
- (3) (I)-(C), (II)-(D), (III)-(B), (IV)-(A)
- (4) (I)-(B), (II)-(D), (III)-(A), (IV)-(C)

Official Ans. by NTA (4)

- Sol. (I) $Ca(OH)_2$ is used in white wash
 - (II) NaCl is used in preparation of washing soda

 $2NH_{3} + H_{2}O + CO_{2} \longrightarrow (NH_{4})_{2}CO_{3}$ $(NH_{4})_{2}CO_{3} + H_{2}O + CO_{2} \longrightarrow 2NH_{4}HCO_{3}$ $NH_{4}HCO_{3} + NaCl \longrightarrow NH_{4}Cl + NaHCO_{3}(s)$ $2NaHCO_{3} \xrightarrow{\Delta} Na_{2}CO_{3} + CO_{2} + H_{2}O$

(III) CaSO₄.
$$\frac{1}{2}$$
 H₂O (Plaster of Paris) is used for

making casts of statues

- (IV) CaCO₃ is used as an antacid
- **20.** The increasing order of the boiling points of the major products A, B and C of the following reactions will be :





21. For Freundlich adsorption isotherm, a plot of log (x/m) (y-axis) and log p (x-axis) gives a straight line. The intercept and slope for the line is 0.4771 and 2, respectively. The mass of gas, adsorbed per gram of adsorbent if the initial pressure is 0.04 atm, is _ _ _ _ × 10^{-4g}.

 $(\log 3 = 0.4771)$

Official Ans. by NTA (48.00)

Sol.
$$\frac{x}{m} = KP^{\frac{1}{n}}$$

$$\log\left(\frac{x}{m}\right) = \frac{1}{n}\log P + \log K$$

slope =
$$\frac{1}{n} = 2$$

intercept = $\log K = 0.4771$

$$K = 3$$

mass of gas adsorbed per gm of adsorbent = $\frac{x}{m}$

$$\frac{x}{m} = 3 \times (0.04)^2 = 48 \times 10^{-4}$$

22. A solution of phenol in chloroform when treated with aqueous NaOH gives compound P as a major product. The mass percentage of carbon in P is _____. (to the nearest integer)

(Atomic mass : C = 12; H = 1; O = 16)

Official Ans. by NTA (69.00)

Official Ans. by ALLEN (68.85)



Molecular weight of $C_7H_6O_2 = 122$

%C =
$$\frac{12 \times 7 \times 100}{122}$$
 = 68.85 ≈ 69

23. If the solubility product of AB_2 is 3.20×10^{-11} M³, then the solubility of AB_2 in pure water is _____ × 10⁻⁴ mol L⁻¹. [Assuming that neither kind of ion reacts with water]

Official Ans. by NTA (2.00)

Sol.
$$AB_2(s) \xrightarrow{s} A_{(aq.)}^{+2} + 2B_{(aq.)}^{-1} : K_{sp}$$

 $K_{SP} = S^1 \times (2s)^2 = 4s^3$
 $3.2 \times 10^{-11} = 4 \times S^3$
 $S = 2 \times 10^{-4} \text{ M/L}$

24. The rate of a reaction decreased by 3.555 times when the temperature was changed from 40°C to 30°C. The activation energy (in kJ mol⁻¹) of the reaction is _____.

Take; R=8.314 J mol⁻¹ K⁻¹ In 3.555 = 1.268

Official Ans. by NTA (100.00)

Official Ans. by ALLEN (99.98)

Sol.
$$\ell n \left(\frac{K_{T_2}}{K_{T_1}} \right) = \frac{E_a}{R} \left[\frac{1}{T_1} - \frac{1}{T_2} \right]$$

$$\Gamma_1 = 303 \text{ K}$$
; $T_2 = 313 \text{ K}$

$$\frac{K_{T_2}}{K_{T_1}} = 3.555$$

$$\ell n(3.555) = \frac{E_a}{8.314} \left[\frac{1}{303} - \frac{1}{313} \right]$$

$$E_a = 99980.715$$

$$E_a = 99.98 \frac{kJ}{mole}$$

25. The atomic number of Unnilunium is _____.

Official Ans. by NTA (101.00)

Sol. Unnilunium $\Rightarrow 101$